Serial No.: 10/789,081 : February 27, 2004

Page : Page 2 of 22

Amendments to the Claims:

The following listing replaces all previous listings of the claims:

1. (Currently amended) A probe array for qualitative and/or quantitative detection of target molecules in a sample by molecular interactions between probe molecules and target

molecules on the probe array, comprising

an array surface.

a first cleavage product of a first probe molecule, wherein the first cleavage product of the first probe molecule is bound to a first region of a target molecule, and wherein the first cleavage product of the first probe molecule includes a label:

a second cleavage product of the first probe molecule immobilized on the array surface at a first defined site, wherein the second cleavage product of the first probe molecule is bound to a second region of the target molecule; and

a cleavage product of a second probe molecule immobilized on the array surface at a second defined site, wherein the cleavage product of the second probe molecule is not bound to a target molecule; wherein the cleavage products of the first and second probe molecules are in contact with a cleaving solution.

- 2. (Previously presented) The probe array of claim 1, wherein the first and second probe molecules are selected from the group consisting of oligonucleotides, peptides, proteins and their analogues.
- 3. (Previously presented) The probe array of claim 1, wherein the first and second probe molecules are oligonucleotides.
 - 4. (Original) The probe array of claim 3, wherein the oligonucleotides have a length of

Serial No.: 10/789,081 Filed: February 27, 2004 Page: Page 3 of 22

from 10 to 100 bases

(Previously presented) The probe array of claim 1, wherein the first cleavage product of the first probe molecule and the second cleavage product of the first probe molecule are

approximately equal in size.

6. (Previously presented) The probe array of claim 1, wherein the cleavage products of

the first and second probe molecules are products of non-enzymatic cleavage.

7. (Currently amended) The probe array of claim 1, wherein the cleavage products of [[he]] the first and second probe molecules are products of cleavage by chemical and/or physical

methods.

 (Previously presented) The probe array of claim 1, wherein the cleavage products of the first and second probe molecules are products of cleavage by acid anions, base cations,

fluoride and/or heavy metal ions.

9. (Original) The probe array of claim 8, wherein the heavy metal ions comprise

mercury ions and/or silver ions.

10. (Previously presented) The probe array of claim 1, wherein cleavage products of the

first and second probe molecules are products of cleavage by photolysis.

11. (Previously presented) The probe array of claim 1, wherein the cleavage products of the first and second probe molecules are products of cleavage of a nucleic acid of the formula

A₁-S-A₂, wherein S is a nucleic acid that comprises the at least one selectively cleavable bond,

and A1 and A2 are any nucleic acids or nucleic acid analogues.

Attorney's Docket No.: 15111.0080 / CLON0010

Applicant : Ellinger et al.

Serial No. : 10/789,081

Filed : February 27, 2004

Page : Page 4 of 22

12. (Original) The probe array of claim 11, wherein S is a nucleotide dimer that is bridged by the selectively cleavable bond.

13. (Currently amended) The probe array of claim 12, wherein S is selected from the group consisting of the following dimers having the formulae I and II:

wherein X and Y are independently selected from the group consisting of O, NH and S, provided that X and Y are not simultaneously O; and B represents a nucleobase which is adenine, guanine, cytosine or thymine,

[[a)]]

[[b)]]

II

Serial No.: 10/789,081 Filed: February 27, 2004 Page: Page 5 of 22

wherein X and Y are independently selected from the group consisting of O, NH and S, provided that X and Y are not simultaneously O, if PG is not a labile protective group; B represents a nucleobase which is adenine, guanine, cytosine or uracil; and PG is selected from the group consisting of H and labile protective groups.

- 14. (Previously presented) The probe array of claim 1, wherein the cleavage products of the first and second probe molecules are products of cleavage of a phosphothioate bond.
- 15. (Original) The probe array of claim 1, wherein the label is a detectable unit, which is selected from the group consisting of fluorescent labels, luminescent labels, metal labels, enzyme labels, radioactive labels, polymeric labels and nucleic acids, which are detectable by hybridisation with a labelled reporter probe.
- 16. (Original) The probe array of claim 15, wherein the detectable unit is coupled to the probe molecules via an anchor group.
- 17. (Previously presented) The probe array of claim 1, wherein said array further comprises third probe molecules arranged on at least one array element of the probe array, wherein the third probe molecules have at least one label and no selectively cleavable bond.
- 18. (Previously presented) The probe array of claim 17, wherein the third probe molecules are oligonucleotides having a defined or randomised sequence.
- 19. (Original) The probe array of claim 1, further comprising an array element having arranged thereon detectable units that are not linked to a probe molecule.
- 20. (Previously presented) The probe array of claim 17, wherein the third probe molecules are arranged on different array elements which differ in their labelling degree.

Serial No. : 10/789,081 Filed : February 27, 2004 Page : Page 6 of 22

 (Original) The probe array of claim 19, wherein the detectable units are arranged on different array elements which differ in their labelling degree.

- 22. (Currently amended) The probe array of claim 1, further comprising fourth probe molecules which have no affinity or at least no specific affinity to target molecules molecules, wherein the fourth probe molecules are arranged on at least one array element.
- (Previously presented) The probe array of claim 22, wherein the fourth probe molecules are oligonucleotides with a defined or randomised sequence.
- 24. (Previously presented) The probe array of claim 1, further comprising fifth probe molecules arranged on at least one array element, and which have a specific affinity to spiking target molecules which are externally added to the sample.
- 25. (Currently amended) The probe array of claim 24, comprising array elements distributed over the entire surface of the array, on which said fifth probe molecules are arranged, which have a label and a selectively cleavable bond located between the label and the immobilization site of the probe on the surface and which have a specific affinity to spiking target molecules added externally to the sample or to a target molecule present in the sample in sufficient concentration to lead to a clearly detectable signal.

26.-51. (Canceled)

52. (Original) A kit for qualitative and/or quantitative detection of target molecules from a sample by molecular interactions between probe molecules and target molecules on probe arrays, comprising: a) the probe array of claim 1; b) reagents for the selective cleavage of the selectively cleavable bond in the probe molecules; c) hybridisation buffer; and d) optionally,

Serial No.: 10/789.081 Filed : February 27, 2004

: Page 7 of 22 Page

washing buffer.

53. (Original) The kit of claim 52, wherein the reagents are selected from the group

consisting of heavy metal ions and enzymes.

54. (Original) The kit of claim 53, wherein the heavy metal ions are selected from

mercury ions and/or silver ions.

55. (Original) The kit of claim 52, further comprising a reaction chamber.

56. (Original) The kit of claim 52, further comprising a detection device.

57. (Original) The kit of claim 52, further comprising a temperature control unit.

58. (Original) The kit of claim 52, wherein the probe array is in the form of a highly

integrated autonomous unit.

59.-61. (Canceled)

62. (Previously presented) A probe array, comprising:

an array surface.

a first probe molecule immobilized on the array surface having at least one label and at

least one selectively cleavable bond between the site of immobilization on the array surface and

the label, wherein the first probe molecule is bound to a corresponding target molecule; and a second probe molecule immobilized on the array surface having at least one label and at

least one selectively cleavable bond between the site of immobilization on the array surface and

the label, wherein the second probe molecule is not bound to a corresponding target molecule;

wherein the first and second probe molecules are in contact with a cleaving solution.

Serial No.: 10/789,081 Filed: February 27, 2004 Page: Page 8 of 22

63. (Previously presented) The probe array of claim 62, wherein the first and second probe molecules are selected from the group consisting of oligonucleotides, peptides, proteins and their analogues.

- 64. (Previously presented) The probe array of claim 62, wherein the first and second probe molecules are oligonucleotides.
- 65. (Previously presented) The probe array of claim 64, wherein the oligonucleotides have a length of from 10 to 100 bases.
- 66. (Previously presented) The probe array of claim 62, wherein the selectively cleavable bond of the first probe molecule is located approximately in the centre between the site of the immobilization of the probe molecule on the array surface and the label.
- 67. (Previously presented) The probe array of claim 62, wherein the selectively cleavable bond of the first probe molecule cannot be selectively cleaved by enzymatic methods.
- 68. (Previously presented) The probe array of claim 62, wherein the selectively cleavable bond of the first probe molecule can be selectively cleaved by chemical and/or physical methods.
- 69. (Previously presented) The probe array of claim 62, wherein the selectively cleavable bond of the first probe molecule can be selectively cleaved by the addition of acid anions, base cations, fluoride and/or heavy metal ions.
- 70. (Previously presented) The probe array of claim 69, wherein the heavy metal ions comprise mercury ions and/or silver ions.

Serial No. : 10/789,081 Filed : February 27, 2004 Page : Page 9 of 22

71. (Previously presented) The probe array of claim 62, wherein the selectively cleavable bond of the first probe molecule can be selectively cleaved by photolysis.

- 72. (Previously presented) The probe array of claim 62, wherein the first probe molecule comprises a nucleic acid of the formula A₁-S-A₂, wherein S is a nucleic acid that comprises the at least one selectively cleavable bond, and A₁ and A₂ are any nucleic acids or nucleic acid analogues.
- 73. (Previously presented) The probe array of claim 72, wherein S is a nucleotide dimer that is bridged by the selectively cleavable bond.
- 74. (Previously presented) The probe array of claim 73, wherein S is selected from the group consisting of the following dimers having the formulae I and II:

wherein X and Y are independently selected from the group consisting of O, NH and S, provided that X and Y are not simultaneously O; and B represents a nucleobase which is adenine, guanine, cytosine or thymine,

Serial No. : 10/789,081 Filed : February 27, 2004 Page : Page 10 of 22

wherein X and Y are independently selected from the group consisting of O, NH and S, provided that X and Y are not simultaneously O, if PG is not a labile protective group; B represents a nucleobase which is adenine, guanine, cytosine or uracil; and PG is selected from the group consisting of H and labile protective groups.

- 75. (Previously presented) The probe array of claim 62, wherein the selectively cleavable bond of the first probe molecule is a phosphothicate bond.
- 76. (Previously presented) The probe array of claim 62, wherein the label is a detectable unit, which is selected from the group consisting of fluorescent labels, luminescent labels, metal labels, enzyme labels, radioactive labels, polymeric labels and nucleic acids, which are detectable by hybridisation with a labelled reporter probe.
- 77. (Previously presented) The probe array of claim 76, wherein the detectable unit is coupled to the probe molecules via an anchor group.
- 78. (Previously presented) The probe array of claim 62, wherein said array further comprises third probe molecules arranged on at least one array element of the probe array, wherein the third probe molecules have at least one label and no selectively cleavable bond.

Serial No.: 10/789,081 Filed: February 27, 2004 Page: Page 11 of 22

79. (Previously presented) The probe array of claim 78, wherein the third probe molecules are oligonucleotides having a defined or randomised sequence.

- 80. (Previously presented) The probe array of claim 62, further comprising an array element having arranged thereon detectable units that are not linked to a probe molecule.
- 81. (Previously presented) The probe array of claim 78, wherein the third probe molecules are arranged on different array elements which differ in their labelling degree.
- 82. (Previously presented) The probe array of claim 80, wherein the detectable units are arranged on different array elements which differ in their labelling degree.
- 83. (Previously presented) The probe array of claim 62, further comprising fourth probe molecules which have no affinity or at least no specific affinity to target molecules, wherein the fourth probe molecules are arranged on at least one array element.
- 84. (Previously presented) The probe array of claim 83, wherein the fourth probe molecules are oligonucleotides with a defined or randomised sequence.
- 85. (Previously presented) The probe array of claim 62, further comprising fifth probe molecules arranged on at least one array element, and which have a specific affinity to spiking target molecules which are externally added to the sample.
- 86. (Previously presented) The probe array of claim 85, comprising array elements distributed over the entire surface of the array, on which said fifth probe molecules are arranged, which have a label and a selectively cleavable bond located between the label and the immobilization site of the probe on the surface and which have a specific affinity to spiking

Applicant : Ellinger et al. Attorney's Docket No.: 15111.0080 / CLON0010 Serial No. : 10/789,081

Serial No. : 10/789,081 Filed : February 27, 2004 Page : Page 12 of 22

target molecule added externally to the sample or to a target molecule present in the sample in sufficient concentration to lead to a clearly detectable signal.

87.-89 (Canceled)